



Received: 22.1.2022

Revised: 25.6.2022

Accepted: 1.7.2022

Published: 8.7.2022

Potravinárstvo Slovak Journal of Food Sciences

vol. 16, 2022, p. 367-374

<https://doi.org/10.5219/1735>

ISSN: 1337-0960 online

[www.potravinarstvo.com](http://www.potravinarstvo.com)

© 2022 Authors, CC BY 4.0

## Incorporation of catechin extracts from gambier products and pasak bumi in the production of functional instant green robusta coffee

*Budi Santoso, Muhammad Ridho Wahyu Aulia, Syerina Raihatul Jannah, Gatot Priyanto, Agus Wijaya, Hermanto Hermanto*

### ABSTRACT

The research was used to produce functional instant green coffee through gambier catechin extract and pasak bumi powder. This involved using a non-factorial completely randomized design with 5 treatments and 3 replications. The treatments consist of 5 formulations (F), including the instant green coffee (%), gambier catechin extract (%), and pasak bumi powder (%) where F1 was at 100:0:0, F2 was 80:15:5, F3 was 70:20:10, F4 was 60:25:15, and F5 was 50:30:20. The results showed the functional instant green coffee produced has a water content of 3.84 – 4.81%, soluble speed of 26.78 – 29.33 seconds, and total phenol of 16.79 – 169.48 mg/L, and IC<sub>50</sub> of 44.68 – 207.59 ppm. The addition of gambier catechin extract and pasak bumi powder to the formulation was observed to have significantly increased the functional properties and water content. Moreover, the soluble speed of the instant coffee fulfils the quality requirements of the Indonesian National Standard (SNI) number 2983 of 2014.

**Keywords:** gambier, instant, catechin, green coffee, pasak bumi

### INTRODUCTION

Humans accept coffee from both the sensory and functional aspects despite numerous pieces of information on its effects on body health. It has been reported that both robusta and arabica generally contain functional compounds in chlorogenic acid. This compound was also discovered by [34] to be present in coffee as an antioxidant, with robusta reported by [38] to contain higher content at 43.63% than arabica, which has 36.18%. According to [14], roasting can reduce robusta caffeine and chlorogenic acid levels by 13 – 25% and 37 – 59%, respectively. Several studies have been conducted to maintain the antioxidant properties of coffee such as the addition of herbal cereals in [29], optimisation of roasting temperature to reduce damage to chlorogenic acid compounds in [8] and [3], and the use of a spontaneous fermentation with *Wickerhamomyces anomalus* (Strain KNU18Y3) on green coffee beans in [7].

Green coffee is currently gaining popularity among world coffee lovers, and it is mainly different from the ordinary types due to the effect of the bean processing method on its functional properties and aroma. According to [6], green robusta has better functional properties than roasted coffee, as indicated by their total phenol contents of 208.89 mg/L and 119.22 mg/L, respectively. [18] also showed that green robusta contains 81.6% antioxidant compounds and has higher caffeine content and high antioxidant properties. It is important to add bioactive compound materials in its production process to increase its antioxidant properties and reduce caffeine levels. One source of these bioactive compounds is catechin and pasak bumi extract.

Catechin is a product from the aqueous extraction of the leaves and twigs of the gambier plant (*Uncaria gambir Roxb*), which have been discovered to contain more than 52.25% catechin compounds [37]. This extract was further reported by [11] to be an antioxidant with an IC<sub>50</sub> of 2.74 g/mL, while [30] also showed its ability to form canna-based edible films, which are antioxidants. According to [13] and [36], the roots of the pasak bumi plant also contain eurikomanone, quassinoids, flavonoid, phenolic, and terpenoid compounds which are observed to have antioxidant potentials.

### Scientific Hypothesis

The addition of gambir catechin extract has a significant effect on increasing the functional properties of instant green coffee, especially its antioxidant activity.

## MATERIAL AND METHODOLOGY

### Samples

Instant coffee powder made from green robusta coffee powder incorporated with gambir catechin extract.

### Chemicals

The materials used consist of distilled water, tannic acid, 96% ethanol, 2,2-diphenyl-1-picrylhydrazil (DPPH), folin-ciocalteu, methanol, Na<sub>2</sub>CO<sub>3</sub>, and nutrient broth (NB) obtained from the Laboratory of Chemical Agricultural Products, Faculty of Agriculture, Sriwijaya University, Indonesia.

### Biological Material

Gambier powder from Babat Toman Village, Musi Banyuasin Regency, South Sumatra, Indonesia. Robusta green coffee powder from JagadRaye Coffee micro and small enterprise in Pagar Alam, South Sumatra, Indonesia. Pasak bumi powder from the Laboratory of Chemical Agricultural Products, Faculty of Agriculture, Sriwijaya University, Indonesia.

### Instruments

The tools used include an autoclave, blender (Philips, Holland), hot plate, incubator (Memmert, Germany), filter paper, laminar airflow (LAF), brand analytical balance (Kenko, Japan), drying oven (Memmert, Germany), pH meter (Eutech, Malaysia), micropipette (Dragon Lab, China), rotary vacuum evaporator, 80 mesh filter, spectrophotometer (A and E Lab, USA), and vortex (Digisystem, Taiwan).

### Laboratory Methods

The parameters evaluated include water content [2]: measurement of water content using the gravimetric method. Soluble speed [2]: Dissolve 100 g of instant coffee in 200 mL of water. Then the length of time instant coffee dissolves in water is calculated as the speed at which it dissolves in water using a stopwatch. Total phenol [31]: Determination of total phenol content was carried out by means of a spectrophotometric method using Folin-Ciocalteu reagent. Antioxidant activity [17]: Antioxidant testing using the DPPH method (2,2 diphenyl-1-picrylhydrazyl) was used.

### Description of the Experiment

**Sample preparation:** The instant green coffee powder, gambier catechin extract, and instant pasak bumi powder with a size of 80 mesh are mixed. Each treatment is put into a cup and then brewed with 100 mL of hot water at 80 °C and stirred using a magnetic stirrer.

**Number of samples analyzed:** A non-factorial completely randomized design was used in this study. A total of five treatments are carried out using the percentage ratio of instant green coffee: gambier product catechin extract: instant pasak bumi. F1 = (100:0:0), F2 = (80:15:5), F3 = (70:20:10), F4 = (60:25:15), and F5 = (50:30:20).

**Number of repeated analyses:** Three repetitions for each treatment factor. The total sample analysed was 15 samples.

**Number of experiment replication:** Each treatment was repeated 3 times.

### Design of the experiment:

#### Instant green coffee

Green coffee beans were dried to a moisture content of 12% and ground using a grinder. The powder was filtered using an 80-mesh sieve, after which water was added at a temperature of 100 °C and a ratio of 1:2, stirred, left for 10 minutes, and later filtered using a filter cloth to obtain the filtrate. Moreover, maltodextrin (10% w/w) and egg white (20% w/w) were added to the filtrate, mixed using a mixer for 10 minutes at high speed to form foam, and spread out on an aluminium pan lined with Polypropylene plastic. The mixture was dried in a carbine dryer at 60 °C for 4 hours, blended, and filtered using an 80-mesh filter to obtain a green coffee powder.

#### Gambier product catechin extract

The catechin extract was prepared using the maceration method. This involved blending the dried gambier sticks until smooth and sieved through an 80-mesh sieve. The 100g gambier powder was macerated using ethanol for 1 day (24 hours) at a ratio of 3:1. Moreover, the catechin extract was filtered using Whatman filter paper No. 41 and evaporated at 85 °C with a rotary vacuum evaporator to vaporise the ethanol and remove the aroma. The catechin extract was later dried using an oven at a temperature of 85 °C for approximately 20 hours, blended, and sifted again.

#### Instant pasak bumi powder production

The instant pasak bumi powder was prepared. This involved the filtration of the powder using an 80-mesh sieve after which water was added at 1:2 and a temperature of 100 °C; the mixture was stirred, left for 10 minutes, and filtered again using a filter cloth to obtain the pasak bumi filtrate. Moreover, maltodextrin (10% w/w) and egg

white (20% w/w) were added to the filtrate, mixed using a mixer for 10 minutes at high speed to form foam, and spread out on an aluminium pan lined with Polypropylene plastic. The mixture was dried in a carbine dryer at a temperature of 60 °C for 4 hours, blended, and filtered using an 80-mesh filter to obtain a green coffee powder.


### Statistical Analysis

This study used a factorial completely randomized design. The treatment with a significant effect was further tested using the honest real difference test (HSD) at = 5%. The data were analysed using the SAS software version of Windows 9 to analyse of variance.

## RESULTS AND DISCUSSION

### Water content

The water content of the functional instant green coffee produced ranged from 3.84 to 4.81% with the highest and lowest recorded in F5 and F1 treatments respectively as indicated in the following Figure 1.



**Figure 1** Effect of formulation on the water content of functional instant green coffee. Note: F1 = 100% green coffee instant: 0% gambir catechin extract: 0% instant pasak bumi; F2 = 80% green coffee instant: 15% gambir catechin extract: 5% instant pasak bumi; F3 = 70% green coffee instant: 20% gambir catechin extract: 10% instant pasak bumi; F4 = 60% green coffee instant: 25% gambir catechin extract: 15% instant pasak bumi; F5 = 50% green coffee instant : 30% gambir catechin extract: 20% instant pasak bumi.

The diversity analysis in Figure 1 showed that the formulation treatment significantly affects the water content of functional instant green coffee. Moreover, the F3 treatment with 20% gambir catechin extract and 10% pasak bumi was observed to have increased the water content. This is associated with the fact that the catechin extract and pasak bumi contain phenolic compounds with a hydroxyl group (OH) that can bind water. It is also important to note that the existence of more OH groups usually leads to more water being bound. Meanwhile, the water content in foodstuffs comprises both bound and free water.

This instant coffee fulfils the quality requirements of the Indonesian National Standard (SNI) No. 2983 of 2014 which states that the maximum water content is 5%. The values obtained in this research were observed to be higher than the 1.57 – 1.61% reported by [21] for instant coffee from Tungkal Jambi and the 2.34% by [39] for cold-brewed instant coffee. Meanwhile, the values are in the same range as 4.4.% found by [15] for instant coffee produced from micro-size coffee combined with *Bacillus coagulans*.

### Soluble Speed

This is one of the quality requirements for instant coffee according to SNI No. 2983 of 2014, which is set at a maximum of 30 seconds. The values obtained in this research were between 26.78 – 29.33 seconds, as indicated in Figure 2 and this means the requirements are satisfied. Meanwhile, the values are higher than the 152.26 seconds [19] for instant coffee made from robusta coffee incorporating maltodextrin but lower than the 11.48 – 13.95 seconds reported by [28] while studying instant robusta with coconut sugar and cane sugar.

The diversity analysis showed that the formulation treatment significantly affects the soluble speed of functional instant green coffee. A higher concentration of gambier catechin extract in the formulation was found to cause a reduction in the soluble speed as indicated in Figure 2. This is because the catechin compounds in gambier products are semi-polar and a higher concentration of catechin usually leads to higher semi-polar nature of instant coffee, thereby, causing a reduction in the solubility of the product in water. This phenomenon was also reported in [24].




Figure 2 Effect of formulation treatment on the soluble speed of functional instant green coffee.

**Total Phenol**

The total phenol of the functional instant green coffee produced ranged from 16.79 to 169.48 mg/L as indicated in Figure 3.




Figure 3 Effect of formulation treatment on total phenol of functional instant green coffee.

These values are slightly lower than 171.633 mg/L reported by [4] and higher than 16.26 – 30.65 mg/L and 42.4 – 59.8 mg/L recorded by [32], [33] and [5], respectively. However, this coffee has a total phenol content similar to the results of research by [9], which is 29.23 – 158.19 mg/mLGAE, [22] regarding cinnamon coffee of 34.46 mg/mLGAE, oven-roasted coffee, which is 16 – 66 mg/mLGAE [1], famous brand coffee circulating in Indonesia is 46.27 mg/mLGAE [16] and roasted arabica coffee is 49.90 mg/mLGAE [23]. Compared with the research of [6], this total phenol is much lower, i.e. unroasted coffee contains 208.89 mg/mLGAE of total phenol and 119.22 mg/mLGAE in roasted coffee.


The diversity analysis showed the significant effect of the formulation treatment on the total phenol of functional instant green coffee. It was discovered that a higher concentration of gambier catechin extract and pasak bumi in

the formulation increased the total phenol. This is, therefore, associated with the polyphenolic compounds in the catechin extract and pasak bumi. The result is in line with the findings of [20] and [29] that gambier contains polyphenol compounds in the form of catechins by 50%. In comparison, [40] found phenolic compounds of catechins and tannins at 65.6 – 74.2% and 11.32 – 17.76%, respectively. Moreover, [10] showed that pasak bumi contains several secondary metabolites: alkaloids, terpenoids, sterpenoids, steroids, flavonoids (phenols), and saponins.

### Antioxidant Activity

The antioxidant activity of functional instant green coffee was measured using  $IC_{50}$  such that a higher  $IC_{50}$  value indicates lower antioxidant activity and vice versa. The values were observed to be from 44.68 – 207.59 ppm as shown in Figure 4, and are the same as the findings of [25] that the encapsulated green coffee extract has 87.65 ppm and [39], which showed that green coffee brewed with cold water has 71.97 – 83.21 ppm. However, the values are higher than the 25.187 ppm reported for green coffee extract dried using the foam mat method by [26], [27], and [18] reported that robusta green coffee contains antioxidants with an  $IC_{50}$  of 81.6  $\mu\text{g/mL}$  and lower than 167.426 to 294.710 ppm recorded for green coffee from Ethiopia by [35] and [12] reported that robusta coffee contains antioxidants with an  $IC_{50}$  of 2210  $\mu\text{g/mL}$ .

The diversity analysis showed that the formulation treatment significantly affects the  $IC_{50}$  of functional instant green coffee, as indicated in Figure 4. This was observed because a higher concentration of gambier catechin extract and pasak bumi powder in the formulation caused a reduction in the  $IC_{50}$  and a higher antioxidant activity. This is associated with flavonoid compounds that are considered antioxidants in the gambier catechin extracts and pasak bumi powder. Moreover, it also indicates consistency with the total phenol data recorded in Figure 3, which showed the same trend. Phenol is also an antioxidant, which means a higher content of this compound can increase the antioxidant properties of the product, as indicated by a decrease in  $IC_{50}$ .



**Figure 4** Effect of formulation treatment on  $IC_{50}$  of functional instant green coffee.

### CONCLUSION

Added catechin extract of gambier and pasak bumi in instant green coffee significantly increases total phenol content and  $IC_{50}$ . Besides that, there was also a change in the physical properties of instant green coffee, namely an increase in water content and speed of dissolving. The functional instant green coffee produced has a water content value of 3.84 – 4.81%, soluble speed of 26.78 – 29.33 s, total phenol of 16.79 – 169.48 mg/L and an  $IC_{50}$  of 44.68 – 207.59 ppm.

### REFERENCE

1. Saeed Alkaltham, M., Musa Özcan, M., Uslu, N., Salamatullah, A. M., & Hayat, K. (2020). Effect of microwave and oven roasting methods on total phenol, antioxidant activity, phenolic compounds, and fatty acid compositions of coffee beans. In *Journal of Food Processing and Preservation* (Vol. 44, Issue 11). Wiley. <https://doi.org/10.1111/jfpp.14874>
2. AOAC. 2012. *Official Methods of Analysis* (18th edition) Association of Official Analytical Chemists International. Maryland, USA.
3. Bobková, A., Hudáček, M., Jakobová, S., Belej, L., Capcarová, M., Čurlej, J., Bobko, M., Árvay, J., Jakab, I., Čapla, J., & Demianová, A. (2020). The effect of roasting on the total polyphenols and antioxidant activity

- of coffee. In *Journal of Environmental Science and Health, Part B* (Vol. 55, Issue 5, pp. 495–500). Informa UK Limited. <https://doi.org/10.1080/03601234.2020.1724660>
4. Christianty, F. M., Holidah, D., Fajrin, F. A., Salsabina, M. C. A., & Roni, A. (2020). The Lipid Profile and Aorta Histopathology on Hyperlipidemic Rat by Giving Green Coffee Extract. In *Indonesian Journal of Pharmaceutical Sciences* (Vol. 18, Issue 1, pp. 21–27). Universitas Gadjah Mada - Faculty of Pharmacy.
  5. Cheng, K., Dong, W., Long, Y., Zhao, J., Hu, R., Zhang, Y., & Zhu, K. (2019). Evaluation of the impact of different drying methods on the phenolic compounds, antioxidant activity, and in vitro digestion of green coffee beans. In *Food Science & Nutrition* (Vol. 7, Issue 3, pp. 1084–1095). Wiley. <https://doi.org/10.1002/fsn3.948>
  6. Górnaś, P., Dwiecki, K., Siger, A., Tomaszewska-Gras, J., Michalak, M., & Polewski, K. (2015). Contribution of phenolic acids isolated from green and roasted boiled-type coffee brews to total coffee antioxidant capacity. In *European Food Research and Technology* (Vol. 242, Issue 5, pp. 641–653). Springer Science and Business Media LLC. <https://doi.org/10.1007/s00217-015-2572-1>
  7. Haile, M., & Kang, W. H. (2020). Antioxidant Properties of Fermented Green Coffee Beans with *Wickerhamomyces anomalus* (Strain KNU18Y3). In *Fermentation* (Vol. 6, Issue 1, p. 18). MDPI AG. <https://doi.org/10.3390/fermentation6010018>
  8. Herawati, D., Giriwono, P. E., Dewi, F. N. A., Kashiwagi, T., & Andarwulan, N. (2018). Critical roasting level determines bioactive content and antioxidant activity of Robusta coffee beans. In *Food Science and Biotechnology* (Vol. 28, Issue 1, pp. 7–14). Springer Science and Business Media LLC. <https://doi.org/10.1007/s10068-018-0442-x>
  9. Ibtisam, K., & Karim. (2013). Optimization of instan controlled pressure drop dic-assisted solvent extraction of total phenols of green coffee beans. In *Journal of Food Studies* (Vol. 2, Issue 1, pp. 42-61). Macrothink Institute.
  10. Iriawati, Rahmawati, A., & Esyanti, R. R. (2014). Analysis of Secondary Metabolite Production in Somatic Embryo of Pasak Bumi (*Eurycoma Longifolia* Jack.). In *Procedia Chemistry* (Vol. 13, pp. 112–118). Elsevier BV. <https://doi.org/10.1016/j.proche.2014.12.014>
  11. Ismail, A. S., Rizal, Y., Armenia, A., & Kasim, A. (2021). Determination of the best method for processing gambier liquid by-product [*Uncaria gambir* (hunter) roxb] as natural antioxidant sources. In *Journal of the Indonesian Tropical Animal Agriculture* (Vol. 46, Issue 2, pp. 166–172). Institute of Research and Community Services Diponegoro University (LPPM UNDIP). <https://doi.org/10.14710/jitaa.46.2.166-172>
  12. Isnindar, Wahyuono, S., & Widyarini, S. (2017). The antioxidant activity of Merapi green coffee berries. In *Journal of Pharmaceutical Science and Clinical Research* (Vol. 2, Issue 2, pp. 130–136).
  13. Khanam, Z., Wen, C. S., & Bhat, I. U. H. (2015). Phytochemical screening and antimicrobial activity of root and stem extracts of wild *Eurycoma longifolia* Jack (Tongkat Ali). In *Journal of King Saud University - Science* (Vol. 27, Issue 1, pp. 23–30). Elsevier BV. <https://doi.org/10.1016/j.jksus.2014.04.006>
  14. Kuncoro, S., Sutiarso, L., Karyadi, J. N. W., & Masithoh, R. E. (2018). Kinetika Reaksi Penurunan Kafein dan Asam Klorogenat Biji Kopi Robusta melalui Pengukusan Sistem Tertutup. In *Agritech* (Vol. 38, Issue 1, p. 105). Universitas Gadjah Mada. <https://doi.org/10.22146/agritech.26469>
  15. Ko, B. S., Lim, S. H., & Han, S. H. (2017). Quality Characteristics of Instant Coffee with Probiotics and Microground Coffee. In *Culinary Science and Hospitality Research* (Vol. 23, Issue 8, pp. 153–162). <https://doi.org/10.20878/cshr.2017.23.8.015>
  16. Lelyana, R. & Cahyono, B. (2015). Total phenolic acid contents in some commercial brands of coffee from Indonesia. In *Journal of Medical Plant and Herbal Therapy Research* (Vol. 3, pp. 27–29). BluePen Journals.
  17. Maesaroh, K., Kurnia, D., & Al Anshori, J. (2018). Perbandingan Metode Uji Aktivitas Antioksidan DPPH, FRAP dan FIC Terhadap Asam Askorbat, Asam Galat dan Kuersetin. In *Chimica et Natura Acta* (Vol. 6, Issue 2, p. 93). Universitas Padjadjaran. <https://doi.org/10.24198/cna.v6.n2.19049>
  18. Masek, A., Latos-Brozio, M., Kałużna-Czaplińska, J., Rosiak, A., & Chrzescijanska, E. (2020). Antioxidant Properties of Green Coffee Extract. In *Forests* (Vol. 11, Issue 5, p. 557). MDPI AG. <https://doi.org/10.3390/fl1050557>
  19. Matarani, F., Mursalin, & Gusriani, I. (2019). The effect of increasing the concentration of maltodextrin on the quality of instant coffee from robusta coffee grounds using a vacuum dryer. In *Proceedings of SEMIRATA BKS-PTN West Region in the Field of Agricultural Science* (Vol. 1, Issue 1, pp. 922-941).
  20. Melia, S., Novia, D., & Juliyarsi, I. (2015). Antioxidant and Antimicrobial Activities of Gambir (*Uncaria gambir* Roxb) Extracts and Their Application in Rendang. In *Pakistan Journal of Nutrition* (Vol. 14, Issue 12, pp. 938–941). Science Alert. <https://doi.org/10.3923/pjn.2015.938.941>
  21. Mursalin, M., Nizori, A., & Rahmayani, I. (2019). The Effect of Heating Schedule on Physico-Chemical Properties of Instant Coffee of Liberika Tungkal Jambi. In *Indonesian Food Science & Technology*



- Journal (Vol. 2, Issue 2, pp. 26–29). Faculty of Education and Teacher Training, Jambi University. <https://doi.org/10.22437/ifstj.v2i2.9442>
22. Nichmah, L., Yuwanti, S., & Suwasono, S. (2019). Kopi kayu manis celup dengan variasi tingkat penyangraian kopi dan konsentrasi bubuk kayu manis. In *Berkala Ilmiah Pertanian* (Vol. 2, Issue 2, p. 50). UPT Penerbitan Universitas Jember. <https://doi.org/10.19184/bip.v2i2.16168>
  23. Odžaković, B., Džinić, N., Kukrić, Z., & Grujić, S. (2016). Effect of roasting degree on the antioxidant activity of different Arabica coffee quality classes. In *Acta Scientiarum Polonorum Technologia Alimentaria* (Vol. 15, Issue 4, pp. 409–417). Uniwersytet Przyrodniczy w Poznaniu (Poznan University of Life Sciences). <https://doi.org/10.17306/j.afs.2016.4.39>
  24. Pambayun, R., Gardjito, M., Sudarmadji, S., & Kuswanto, K. R. (2007). Phenolic content and antibacterial properties of various extracts of gambir (*Uncaria gambir* Roxb). In *Indonesian Journal of Pharmacy* (Vol. 18, Issue 3, pp. 141–146). Faculty of Pharmacy Universitas Gadjah Mada.
  25. Vareltzis, P. K., Zeleskidou, M., Kiroglou, S., & Gargali, I. (2020). Production of instant coffee from cold brewed coffee; process characteristics and optimization. In *Food Science and Applied Biotechnology* (Vol. 3, Issue 1, p. 39). University of Food Technologies - Plovdiv. <https://doi.org/10.30721/fsab2020.v3.i1.92>
  26. Pranowo, D., Adiatmi, A. Y., & Dewi, I. A. (2021). Production optimization of green coffee extracts from Jember robusta (*Coffeacanephora*) coffee using foam mat drying method. In *IOP Conference Series: Earth and Environmental Science* (Vol. 733, Issue 1, p. 012100). IOP Publishing. <https://doi.org/10.1088/1755-1315/733/1/012100>
  27. Pranowo, D., Perdani, C. G., Prihardhini, T. A., Wijana, S., Fahmi, A. S. & Arisandi, D. M. (2020). Optimization of microencapsulation process of green coffee extract with spray drying as a dietary supplement. In *Systematic Reviews in Pharmacy* (Vol. 11, Issue 10, pp. 715–721). Wolters Kluwer Medknow Publications.
  28. Praptiningsih Y. S., Tamtarini, I., & Wijayanti, S. (2012). The properties of coconut sugar instant coffee from various ratios of Arabica Robusta coffee and coconut sugar to granulated sugar. In *Journal of Agrotechnology* (Vol. 6, Issue 1, pp. 70–77). OMICS Publishing Group.
  29. Rahmawati, N., & Wachyuni, A. F. (2013). Kandungan fenolik dan aktivitas antioksidan ekstrak daun gambir kering (*Uncaria gambir* (Hunter) Roxb.). In *Jurnal Indonesian Chimica Acta* (Vol. 4, pp. 1–6). Hasanuddin University.
  30. Samsonowicz, M., Regulaska, E., Karpowicz, D., & Leśniewska, B. (2019). Antioxidant properties of coffee substitutes rich in polyphenols and minerals. In *Food Chemistry* (Vol. 278, pp. 101–109). Elsevier BV. <https://doi.org/10.1016/j.foodchem.2018.11.057>
  31. Santoso, B., Hazirah, R., Priyanto, G., Hermanto, D. & Sugito, D. (2019). Utilization of *Uncaria gambir* Roxb filtrate in the formation of bioactive edible films based on corn starch. In *Food Science and Technology* (Vol. 39, Issue 4, pp. 837–842). FapUNIFESP (SciELO). <https://doi.org/10.1590/fst.06318>
  32. Septiani, Dewi, E. N., & Wijayanti, I. (2007). Antibacterial activity of seagrass extract (*Cymodocea rotundata*) against *Staphylococcus* and *Escherichia coli* bacteria. In *Fisheries Saintek* (Vol.13, Issue 1, pp. 1–6).
  33. SNI 2983 : 2014. Instant coffee. National Standardization Board. Jakarta.
  34. Siva, R. (2016). Assessment Of Antioxidant Activity And Total Phenolic Content From Green Coffee Robusta Sp. BEANS. In *Malaysian Journal of Analytical Science* (Vol. 20, Issue 5, pp. 1059–1065). Penerbit Universiti Kebangsaan Malaysia (UKM Press). <https://doi.org/10.17576/mjas-2016-2005-10>
  35. Jeszka-Skowron, M., Frankowski, R., & Zgoła-Grześkowiak, A. (2020). Comparison of methylxantines, trigonelline, nicotinic acid and nicotinamide contents in brews of green and processed Arabica and Robusta coffee beans – Influence of steaming, decaffeination and roasting processes on coffee beans. In *LWT* (Vol. 125, p. 109344). Elsevier BV. <https://doi.org/10.1016/j.lwt.2020.109344>
  36. Tasew, T., Mekonnen, Y., Gelana, T., Redi-Abshiro, M., Chandravanshi, B. S., Ele, E., Mohammed, A. M., & Mamo, H. (2020). In Vitro Antibacterial and Antioxidant Activities of Roasted and Green Coffee Beans Originating from Different Regions of Ethiopia. In *International Journal of Food Science* (Vol. 2020, pp. 1–8). Hindawi Limited. <https://doi.org/10.1155/2020/8490492>
  37. Triawanti, Sanyoto, D. D., & Noor, M. S. (2020). The supplementation of pasak bumi (*Eurycoma longifolia* Jack.) in undernourished rats to increase spatial memory through antioxidant mechanism. In *Clinical Nutrition Experimental* (Vol. 33, pp. 49–59). Elsevier BV. <https://doi.org/10.1016/j.clnex.2020.08.002>
  38. Widiyarti, G., A. Sundowo, E. Filaila, and J.A. Laksmono. (2020). The mechanically extraction process from leaves and twigs of gambir (*Uncaria gambir* Roxb) and its antioxidant activity. In *Journal of Pure and Applied Chemistry Research* (Vol. 9, Issue 1, pp. 18–15). <https://doi.org/10.21776/jpacr.ub.2020.009.01.509>
  39. Wolska, J., Janda, K., Jakubczyk, K., Szymkowiak, M., Chlubek, D., & Gutowska, I. (2017). Levels of

Antioxidant Activity and Fluoride Content in Coffee Infusions of Arabica, Robusta and Green Coffee Beans in According to their Brewing Methods. In *Biological Trace Element Research* (Vol. 179, Issue 2, pp. 327–333). Springer Science and Business Media LLC. <https://doi.org/10.1007/s12011-017-0963-9>

40. Yeni, G., Syamsu, K., Mardiyati, E., & Muchtar, H. (2017). Determination of the process technology for making pure gambier and standardized catechins from random gambier. In *Industrial Research and Development Journal* (Vol. 7, Issue 1, pp 1–10). Technicals Publications.

### Funds:

This paper is part of the Competitive Research result funded by DIPA budget of Public Service Agency, Sriwijaya University for the fiscal year of 2021, No. 0022/UN9/SK.LP2M.PT/2019, 23 November 2020. with contract No. 0107.045/UN9/SB3.LP2M.PT/2021

### Acknowledgments:

-

### Conflict of Interest:

The authors declare no conflict of interest.

### Ethical Statement:

This article does not contain any studies that would require an ethical statement.

### Contact Address:

\*Budi Santoso, Sriwijaya University: Ogan Ilir, Agricultural Technology Department, Faculty of Agriculture, South Sumatera, Indonesia,

Tel.: +628127853631,

Email: [budisantoso@fp.unsri.ac.id](mailto:budisantoso@fp.unsri.ac.id)

ORCID: <https://orcid.org/0000-0002-5037-0048>

Muhammad Ridho Wahyu Aulia, Sriwijaya University: Ogan Ilir, Agricultural Technology Department, Faculty of Agriculture, South Sumatera, Indonesia,

Tel.: +6281377937776,

Email: [ridho9hspensa@gmail.com](mailto:ridho9hspensa@gmail.com)

ORCID: <https://orcid.org/0000-0002-3051-2635>

Syerina Raihatul Jannah, Sriwijaya University: Ogan Ilir, Agricultural Technology Department, Faculty of Agriculture, South Sumatera, Indonesia,

Tel.: +6282175800458,

Email: [syerinaraihatuljannah@gmail.com](mailto:syerinaraihatuljannah@gmail.com)

ORCID: <https://orcid.org/0000-0002-0989-9222>

Gatot Priyanto, Sriwijaya University: Ogan Ilir, Agricultural Technology Department, Faculty of Agriculture, South Sumatera, Indonesia,

Tel.: +6281233463906,

Email: [tech.gpri@gmail.com](mailto:tech.gpri@gmail.com)

ORCID: <https://orcid.org/0000-0002-0028-5005>

Agus Wijaya, Sriwijaya University: Ogan Ilir, Agricultural Technology Department, Faculty of Agriculture, South Sumatera, Indonesia,

Tel.: +6281377844401,

Email: [agus\\_wijaya@hotmail.com](mailto:agus_wijaya@hotmail.com)

ORCID: <https://orcid.org/0000-0001-8280-2397>

Hermanto Hermanto, Sriwijaya University: Ogan Ilir, Agricultural Technology Department, Faculty of Agriculture, South Sumatera, Indonesia,

Tel.: +6281379133523,

Email: [hermanto.ramlimansyur@gmail.com](mailto:hermanto.ramlimansyur@gmail.com)

ORCID: <https://orcid.org/0000-0002-6926-9767>

Corresponding author: \*

© 2022 Authors. Published by HACCP Consulting in [www.potravinarstvo.com](http://www.potravinarstvo.com) the official website of the *Potravinarstvo Slovak Journal of Food Sciences*, owned and operated by the Association HACCP Consulting, Slovakia, [www.haccp.sk](http://www.haccp.sk). The publisher cooperate with the SLP London, UK, [www.slplondon.org](http://www.slplondon.org) the scientific literature publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License <https://creativecommons.org/licenses/by/4.0>, which permits unrestricted use, distribution, and reproduction in any medium provided the original work is properly cited.