

OPEN ACCESS

Received: 18.3.2023

Revised: 4.5.2023

Accepted: 25.5.2023

Published: 9.6.2023

**Slovak Journal of  
Food Sciences**

*Potravinárstvo Slovak Journal of Food Sciences*

vol. 17, 2023, p. 529-549

<https://doi.org/10.5219/1875>

ISSN: 1337-0960 online

[www.potravinarstvo.com](http://www.potravinarstvo.com)

© 2023 Authors, CC BY-NC-ND 4.0

## **The influence of chitosan on the raspberry quality during the storage process**

*Anastasiia Blahopoluchna, Mikhailo Mushtruk, Nataliia Slobodyanyuk, Nelia Liakhovska, Vladyslav Parakhnenko, Sergiy Udodov, Inna Karpovych, Olena Ochkolyas, Alina Omelian, Gennadiy Rzhevsky*

### **ABSTRACT**

Raspberry is a perishable berry raw material with a high capacity for mechanical and microbiological damage, and therefore, after harvesting, it is necessary to use appropriate technologies to preserve its quality and extend the storage time. This work aimed to study the influence of different concentrations of chitosan solutions on the quality and duration of storage of raspberries under refrigerating conditions. Raspberries were picked at the consumer maturity stage in perforated plastic containers with a capacity of 500 grams. The berries were processed by spraying with 0.5%, 1.0%, and 2.0% chitosan solution, then removing residual moisture. The storage was carried out for twelve days in a refrigerating chamber at a temperature of 2 °C and relative humidity of 95%. The research was performed according to the physical, chemical, and organoleptic indicators, determining the changes in the mass fraction of ascorbic acid, the mass fraction of sugars during storage, taste, aroma, colour, etc. consistency and appearance at the end of storage. It was found that the loss of ascorbic acid in the processed berries was 1.5-3.9 times less than in the reference sample. A similar situation was observed with the mass fraction of sugars, which prevailed 0.9-2.5 times in the processed samples. According to the organoleptic indicators, the samples with a solution concentration of 1.0% and 2.0% were recognized as the best. According to the results of experimental investigations, it was established that pre-processing of berries with chitosan solutions is a promising method to slow down unwanted metabolic processes that take place after harvesting.

**Keywords:** chitosan, storage time, raspberries, pre-processing, biopolymer

### **INTRODUCTION**

Raspberry is a highly nutritious berry crop with a short storage time, high water content and thin cover tissues that mechanical and microbiological factors may easily damage; this berry has a high nutritional value and a wide range of applications in the food industry [1]. Moreover, it can be used to make:

- jams, confitures and other preserves, which have a pleasant taste and aroma, as well as a high pectin content, which allows obtaining a stable gelatinous product;
- fruit juices, nectars and lemonades with a high content of vitamins and antioxidants;
- as an additive to ice cream and yoghurt;
- confectionery and dried fruits.

Raspberries continue their metabolic activity after harvesting, gradually losing quality, decreased resistance to phytopathogenic damage, and increased percentage loss. Therefore, edible coatings made of biopolymers and their combinations extend the storage time of perishable berry products [2].

Chitosan is a biopolymer obtained from chitin, the main component of the skeleton of crustaceans and insects, which can form a thin layer around fresh products, which acts as a protective agent, extending the storage time, and also serves as an inhibitor of metabolic processes [3]. Chitosan has several advantages for processing berries, among which are the following:

- berry storage, chitosan can be used as a preserving agent for berry storage as it is a natural antibacterial agent that helps to prevent the development of spoilage microorganisms [4];
- the storage time extension, chitosan can increase the storage time of berries by creating a protective layer that helps to retain moisture and to prevent oxidation [5];
- the improvement of the quality of berries, chitosan can improve the quality of berries by reducing moisture loss and increasing their viability and resistance to mechanical damage [6];
- the environmental component, chitosan is a natural product that decomposes after use, so it is an environmentally compatible means for raspberry processing [7].

Diseases, improper transport, and storage techniques cost the world economy about \$220 billion annually, reducing crop productivity and quality and leading to higher food prices and global food insecurity. Reducing losses and scraps of fresh fruit and vegetables can help reduce pressure on food production systems, especially in the context of limited natural resources and climate change.

Edible coatings have become integral for protecting fruits and vegetables from phytopathogenic damage. A wide range of studies demonstrated the antimicrobial activity of edible coatings against *Botrytis cinerea*, *Colletotrichum* spp., *Penicillium* spp. and *Alternaria* spp. [8]. Films formed on the surface of fruits can change the atmosphere's composition, creating a barrier for gas exchange, such as oxygen, carbon dioxide and ethylene, which are involved in the process of respiration [9].

There are many edible coatings; among them, chitosan is the safest. Chitosan ( $\beta$ -(1,4)-2-amino-2-deoxy-D-glucose) is a natural biopolymer obtained by deacetylation of chitin, which is the second with ranking most important polysaccharide in nature after cellulose, and which enters into the composition of the exoskeleton structure of marine invertebrates, insects, as well as fungi, algae and yeast [10]. Chitosan is one of the most widely used edible coatings due to its biocompatibility, biodegradability, and bioactivity. When applied to fruits, vegetables and berries, chitosan creates a semi-permeable film that protects against the development of fungus diseases and slows down metabolic processes [11]. This edible coating is widely used for post-harvest fresh fruits and vegetables preservation. The scientific literature on edible coatings using chitosan has increased in recent years. This can be explained by the importance of chitosan for plant protection as a natural fungicide. In 2014, chitosan hydrochloride was approved as one of the first main plant protection substances by the European Union (EU Regulation 2014/563), and the second chitosan formulation was approved in 2022 (EU Regulation 2022/456) [12].

Therefore, the use of raspberry in the food industry is a promising and relevant direction of research because its use will allow to expand of the assortment, increase the quality and nutritional value of products made on its basis, attract new consumers, and export can be a promising line for the development of the food industry of Ukraine.

### Scientific Hypothesis

For further improvement of storing and processing raspberry technology, predicting the storage time depending on various environmental influences may be considered a promising direction. Therefore, the research aimed to develop the most effective method of storing raspberries to improve the prediction of the content of dry soluble substances, sugars and titrating acids in raspberries. By conducting experimental investigations, the expediency of predicting the content of the main components of the chemical composition in raspberries will be determined according to average values, and the factors that have the greatest influence on the accumulation of the studied indicators will be identified.

## MATERIAL AND METHODOLOGY

### Samples

Raspberries of Patricia (Figure 1 a), Polka (Figure 1 b) and Chervona Koroleva (Figure 1 c) varieties were picked in the fields of the academic and research department of the Uman National University of Horticulture at the consumer maturity stage according to DSTU 7179:2010 [13].



a



b



c

**Figure 1** Studied varieties of raspberries: a – berries of the variety of Patricia, b – berries of the variety of Polka, c – berries of the variety of Chervona Koroleva.

## Chemicals

- Acetone, C<sub>3</sub>H<sub>6</sub>O (TD Energobudinvest, Ukraine).
- Sodium hydroxide, NaOH (Khimlaborreaktiv TOV, Ukraine).
- Ascorbic acid, vitamin C (Khimlaborreaktiv TOV, Ukraine).
- Metaphosphoric acid, HPO<sub>3</sub> (Khimlaborreaktiv TOV, Ukraine).
- Pyrocatechin, C<sub>6</sub>H<sub>4</sub>(OH)<sub>2</sub> (Khimlaborreaktiv TOV, Ukraine).
- Chloroform, CHCl<sub>3</sub> (CHEMICO GROUP, Great Britain).
- Methanol, CH<sub>3</sub>OH (CAS, the Netherlands).
- Hexane, C<sub>6</sub>H<sub>14</sub> (Hammerite, the Netherlands).

## Animals, Plants and Biological Materials

For experimental investigations, the following varieties were used: Patricia raspberry variety (produced by the academic and research department of the Uman National University of Horticulture, Cherkasy Oblast, Ukraine); Polka raspberry variety (produced by the academic and research department of the Uman National University of Horticulture, Cherkasy Oblast, Ukraine) and Chervona Koroleva raspberry variety (produced by the academic and research department of the Uman National University of Horticulture, Cherkasy Oblast, Ukraine).

## Instruments

- Drying oven SNOL 67/350 (ThermoEngineering TOV, Ukraine), a titrator (Labor-Technik TOV, Ukraine).
- Analytical electronic balance KERN ABS 120-4 (Khimtex, the State Enterprise, Ukraine).
- Refractometer IRF-454B2M (KOMZ VAT).
- pH meter ULAB MP 511 (ULAB, China).
- Gas chromatograph Kristallux-4000M (Meta-Chrom, the Research and Production Company).
- Refrigerator GGM Gastro (GGM Gastro, Germany).
- Refractometer (IRF-454 B2M, manufactured by Inter-SynteZ TOV, Ukraine).
- Laboratory thermometer (TLS-200, manufactured by Inter-SynteZ TOV, Ukraine).
- Photocolorimeter (KFK-3, manufactured by Inter-SynteZ TOV, Ukraine).
- Flame spectrophotometer (Saturn-4, manufactured by Inter-SynteZ TOV, Ukraine).

## Laboratory Methods

The selection and preparation of samples for analysis of fresh strawberries were carried out according to DSTU (the State Standards of Ukraine) 7205:2009, and the products of its processing – according to DSTU 7244:2009.

The quality assessment of fresh raspberries and preserves was carried out following the following regulatory documents:

- fresh berries - according to DSTU 7205:2009 [14];
- compotes - according to DSTU 8102:2015 [15];
- jams - according to DSTU 4900:2007 [16];

In the test samples, the following parameters were determined:

- an average weight of strawberries by weighing;
- a volume of strawberries by the amount of displaced water when immersed in a measuring cylinder;
- a hardness of berries by calculating the ratio of the mass of berries to their volume;
- a density by piercing a fruit in the equatorial zone with a FT 02 penetrometer;
- a respiration intensity by the amount of released carbon dioxide [17];
- a weight loss of berries by the method of fixed samples weighing [18];
- a shine - visually on a 5-point scale, where 1 – a dull surface of berries, no shine, and 5 - a shiny glossy surface;
- a content of dry soluble substances according to DSTU 8402:2015 [19];
- a content of sugars according to DSTU 4954:2008 [20];
- organic acids by titrating with alkali according to DSTU 4957:2008 [21];
- pH – by the potentiometric method according to DSTU 6045:2008 [22];
- an acetaldehyde content – by the bichromatic and iodometric method [23];
- an ethyl alcohol content – by the iodometric method [24];
- an ascorbic acid content – by the iodometric method [25];
- a content of nitrates – by the ionometric method according to DSTU 4948:2008 [26];

All investigations were performed in triplicate.

The results of the analysis led to the initial mass according to the formula (1):

$$X = \frac{A \times (100 - b)}{100} \quad (1)$$

Where:

$X$  – the content of substances taking into account a mass loss, %;  $A$  – the content of substances at the end of storage, %;  $b$  – the mass loss during the storage period, %.

Microbiological studies were carried out by microscopy using a MICROmede XS – 2610 microscope with a magnification of 50 times, taking samples from the surface of berries with an inoculating wire loop. Photomicrographs were taken using a photo camera.

## Description of the Experiment

**Sample preparation:** Bushes, typical for a certain variety, even-aged, with medium intensity of fruiting, were selected for the research. Raspberries of each variety were picked when the pulp of the berries was still dense enough, but the taste and color were appropriate for this variety. The harvesting date was determined by the following characteristics of the quality of fresh berries: the appearance and the size of berries about the largest longitudinal diameter. The selected berries corresponded to the indicators of the first commercial variety: the shape and colour of the variety, berries with stalks, mechanical damage, vermin damage and fungal diseases. Fruits were picked from different bushes.

Experiment 1: Raspberries were processed with chitosan solutions of six concentrations (0.05%; 0.1%; 0.2%; 0.3%; 0.4%; 0.5%) in two ways: spraying and immersion for 1 min. The processed berries were left to dry completely.

Experiment 2: Raspberries were processed with chitosan solutions of six concentrations: 0.05%; 0.1%; 0.2%; 0.3%; 0.4%; 0.5% and left to dry completely.

Experiment 3: Raspberries were processed with a 0.5% chitosan solution and stored for seven days in a refrigerator with free access to air.

**Number of analyzed samples:** To determine the content of dry soluble substances, sugars and titrating acids, a sample of 100 berries of each pomological variety was taken from 6 bushes that had entered into a full fruiting period.

**Number of repeated analyses:** all instrument measurements and readings were performed 3 times.

**Number of experiment replications:** The number of repeats of each experiment to determine one value was also 3 times.

**Design of the experiment:** Experiment 1: Dry processed berries and the control were weighed and placed in perforated plastic (PET) containers with a capacity of 500 g, and stored at a temperature of 20 °C on racks in a ventilated location. Berries without processing were considered as the control.

Experiment 2: Dry processed berries and the control were weighed and packaged in perforated plastic (PET) containers with a capacity of 500 g and in plastic bags with a thickness of 30 microns. The storage was carried out in two ways: with free access to air and in a modified gas environment at a temperature of 0 ± 2 °C with an atmosphere relative humidity of 90-95%. Berries without processing were considered as the control.

Experiment 3: After storing, the preserves were made from berries according to the manufacturing specifications: "Raspberry puree", "Raspberry jam", "Raspberry confiture", "Raspberry compote", and "Natural raspberry juice". The "Raspberry in own juice" preserves were produced according to the technology developed by our team. The preserves made from fresh unprocessed raw materials were considered as the control.

## Statistical Analysis

The results were evaluated using statistical software Statgraphics Centurion XVII (StatPoint, USA) – multifactor analysis of variance (MANOVA), LSD test. Statistical processing was performed in Microsoft Excel 2016 in combination with XLSTAT. Values were estimated using mean and standard deviations. The reliability of the research results was assessed according to the Student's test at a significance level of  $p \leq 0.05$

## RESULTS AND DISCUSSION

The study was conducted every second day of storage to assess the influence of pre-processing of raspberries with chitosan solutions. We have determined weight loss, a mass fraction of dry soluble substances (a mass fraction of DSS), a respiration intensity, a mass fraction of organic acids, a mass fraction of sugars, an ascorbic acid content, pH level; a density, and a shine degree.

During the storage period ends, the content of ethyl alcohol and acetaldehyde and the yield of commercial products were determined, and the organoleptic evaluation was carried out.


A series of similar scientific investigations, in which the influence of pre-processing with various solutions on various types of fruit and berry raw materials, is described in the following scientific papers:

- the influence of pre-processing of strawberry berries [27], [28];
- the influence of pre-processing of currant berries [29], [30];
- the influence of pre-processing of blackberry berries [31], [32];
- the influence of pre-processing of apple fruits with special solutions [33], [34];
- the influence of pre-processing of apricot fruits with solutions based on sulfites [35], [36];
- the influence of pre-processing of cape gooseberry fruits with special solutions [37], [38].

However, the use of sulfite can cause various types of allergic reactions, so its use is limited.

Raspberries are characterized by a high moisture content, which is lost through thin cover tissues due to quick physiological changes. Berry mass losses during storage are caused by a rather high respiration intensity, a decrease in the content of nutritional substances, and a development of phytopathogenic damage [39].

Pre-processing of raspberry with chitosan solutions made it possible to reduce the weight loss of berries during storage (Figure 2).




**Figure 2** The changes in the natural weight losses of raspberries during refrigerated storage in a modified gas environment (2018-2020).

The mass loss of raspberries increased with each day of storage, and on the second day, it ranged from 0.98 to 1.5% and from 1.0 to 1.3%, depending on the storage method. On the eighth day of the storage, the indicators ranged from 4.3 to 6.3% and from 3.8 to 5.7%. The lowest losses during the entire storage period were detected in the sample with a chitosan processing concentration of 0.5%. During the storage period ends, the mass losses reached 9.6% and 10.9% in control, 6.2-10.4% and 5.2-8.9% in the preprocessed samples.

The analysis of the dynamics of raspberry's weight losses during the two-week storage period demonstrated that the processing with a chitosan solution contributes to weight loss reduction.

Carbohydrates, nitrogenous substances, acids, pectin, vitamins, enzymes, mineral salts and tanning substances represent dry soluble substances. In raspberries, the main part of DSS is carbohydrates, mainly represented by sugars and acids. The change in the mass fraction of dry soluble substances during storage occurs due to the conduction of biochemical processes in raspberries.

The pre-processing of raspberries with chitosan revealed a positive effect on preserving dry soluble substances (Figure 3).




**Figure 3** The change in the mass fraction of dry soluble substances of raspberries during refrigerated storage in a modified gas environment (2018-2020).

The investigations have shown insignificant variations in the mass fraction of dry soluble substances among different methods of refrigerated storage of berries. However, the lowest loss of the mass fraction of dry soluble substances during refrigerated storage with free access to air for 14 days was observed when raspberries were processed with chitosan at a concentration of 0.5%, and the highest loss was in control and at a concentration of 0.05%. A similar dependence of the change in the mass fraction of dry soluble substances was revealed during refrigerated storage in a modified gas environment.

It was proved that the mass fraction of dry soluble substances decreased more slowly in the samples processed with chitosan. On the second day, in the control, the accelerated loss rates of the mass fraction of dry soluble substances were observed, negatively affecting raspberries' preservation.

On the sixth day of the storage of berries with free access to air, an equal value (8.0%) was detected in the sample with a processing concentration of 0.05% and in the sample without any processing. In the future, the difference in indicators between these two samples for both refrigerated storage methods was insignificant (0.2-0.4%), which may indicate the inexpediency of using this concentration. It was found that the lowest changes in the mass fraction of DSS were in samples with a processing concentration of 0.4% and 0.5%.

The respiration intensity is the main indicator of metabolic processes in berries. This is the dominant indicator, the slowing down of which allows for extending the storage time of fruits [40]. Pre-processing raspberries before their placement in storage helps reduce their respiratory activity (Figure 4).



HIP<sub>05</sub> = 2.2

**Figure 4** The changes in the respiration intensity of raspberries during refrigerated storage in a modified gas environment (2018-2020).

The physical effect of chitosan is that a thin transparent film forms on the surface of raspberries, which slows down a gas exchange.

The respiratory exchange of raspberries actively continues after their separation from a parent plant, negatively affecting the storage quality and duration.

The average respiration intensity of fresh raspberries was 34.6 mg CO<sub>2</sub>.kg/h. On the second day of the storage, the indicator decreased sharply regardless of the processing concentration and the storage method and ranged from 8.9 to 10.5 and from 8.5 to 11.3 mg CO<sub>2</sub>.kg/h. This was facilitated by a significant decrease in temperature to 0 ± 2 °C. During further storage, the indicators have continued to decrease gradually. In the sample processed with a 0.5% chitosan solution, the respiration intensity was the lowest and on the eighth day was 5.3 CO<sub>2</sub>.kg/h when stored with free access for air and 4.6 CO<sub>2</sub>.kg/h in a modified gas environment, that by 2.4 and 1.4 less than the control.


Organic acids in raspberries are represented by citric, malic, quinine, salicylic, phosphoric, succinic, shikimic and glycolic acids [41].

During the storage of raspberries, there is a tendency to lose organic acids, which are most involved in respiration. Pre-processing of raspberries with chitosan reduced the respiration intensity, thereby slowing the loss of organic acids by 0.15-0.19% of the counter (Figure 5).

On the second day, the highest percentage of organic acids (0.88%) was found in the sample with a processing concentration of 0.5% for both storage methods.

On the fourth day of the storage, a significant decrease in the content of organic acids was observed in raspberries that were stored with free access to air. However, the samples in a modified gas environment noted a stabilization of the dynamics of losses. On the eighth day of the storage of berries in a refrigerator with free access to air, the lowest content of organic acids (0.50) was detected in the sample with a chitosan processing concentration of 0.1%.





**Figure 5** The change in the mass fraction of organic acids of raspberries during refrigerated storage in a modified gas environment (2018-2020).

At the beginning and until the end of the storage period, the best result was observed in the sample with the highest processing concentration, proving its application's effectiveness.

The processing of berries with chitosan solutions significantly reduces the decomposition rate of organic acids.

It is commonly known that sugars in raspberries are represented by glucose, fructose and sucrose. In combination with organic acids, sugars participate in oxidation processes, so their loss is partly caused by the respiration intensity [42].

Our research established that pre-processing of raspberries with a chitosan solution significantly affects the changes in the content of sugars that occur during the storage period. This is explained by the fact that chitosan slows down the respiratory processes in berries, which causes significant losses of sugars.

It is found that the average sugar content in freshly harvested raspberries was 6.0%. During the entire storage period, significant losses of sugars were detected in the control samples.


The authors of scientific papers [43], [44] found that the average sugar content in freshly harvested raspberries was 6.0-10%, but raspberry varieties and places where these researches have been conducted, were not indicated. Thus, such statements cause doubts about the obtained research results.

On the second day of the storage, a sharp decrease in the mass fraction of sugars was observed in the control samples (5.2%) and (4.9%) and in samples with a chitosan processing concentration of 0.05% (5.0%) and (5.2%). When raspberries were stored in a modified gas environment (MGE) from the beginning to the eighth day, the value for the control and the sample with the minimum processing concentration was equal to one.

By the end of the storage period, the sugar content in berries gradually decreased and, on the fourteenth day, ranged from 2.7 to 4.3% in samples with free access to air and from 2.5 to 3.6% in a modified gas environment.

The intensity of the use of sugars in the physiological and biochemical processes that occurred during the storage of processed raspberries was significantly lower compared to the control due to the slowing down of respiratory activity.

The content of ascorbic acid usually determines the vitamin value of raspberries. The content of vitamin C depends mainly on the variety and soil and climatic conditions.



**Figure 6** The change in ascorbic acid content in raspberries during refrigerated storage with free access to air (2018-2020).


Ascorbic acid is an unstable compound that is easily oxidized during storage [45]. Many factors affect its decomposition; the main of which are light, temperature and pre-processing.

Over the years of investigations, it has been established that the average ascorbic acid content in the Dukat variety raspberries is 65.1 mg/100 g. Depending on the refrigerated storage period, it decreased as well as with free access to air (Figure 6).

It was investigated that the ascorbic acid content in raspberries decreased rapidly, and on the fourth day, it ranged from 50.6 to 58.7 mg/100 g and from 49.5 to 60.2 mg/100 g, depending on the storage method. During the storage period ends, the highest ascorbic acid content was detected in samples with a processing concentration of 0.5% (35.4 mg/100 g) and (37.8 mg/100 g), which is 11.6 and 11.4 more than the control.

The active acidity of berries is an important characteristic of them, as it impacts the microflora's vital activity. The acid taste of fruit and vegetable products is provided by hydrogen ions, which are formed as a result of the electrolytic dissociation of acids and acid salts. The activity of hydrogen ions is characterized by the pH indicator [46].

It was established that the average pH level in fresh raspberries is 3.2. During storage, the active acidity decreases depending on the time and method of storage (Figure 7). Depending on the storage method, the indicator increased by 0.2-0.4 and 0.1-0.3 on the second day. On the eighth day of the storage, the lowest pH level was observed in the sample with a chitosan processing concentration of 0.5% (3.8) and (3.6), 0.5 and 0.4 less than the control.



**Figure 7** The change in the pH level of raspberries during refrigerated storage with free access to air (2018-2020).

During the storage period ends, the acidity of raspberries ranged from 4.2 to 4.8 in samples stored with free access to air and from 4.0 to 4.5 in a modified gas environment.

The analysis of research results proves that chitosan can slow down the pH level increase.

The tissue density indicates the consumer ripeness (degree of maturity) of fruits. It depends on the variety, fruit size and weather conditions during cultivation. The high density contributes to better storing and transporting of fruit and berry raw materials [47]. During storage, the tissue density decreased significantly, and in the middle of the storage period (the 6th day) it ranged from 0.24 to 0.28 kg/cm<sup>2</sup> in processed berries that were under conditions of free access to air, and from 0.25 to 0.28 kg/cm<sup>2</sup> in a modified gas environment.

During ripening, the berry tissue gradually softens, and in the case of frequent rains, it becomes thin and sensitive to mechanical damage.


It was studied that the average density of fresh raspberries after harvesting was 0.30 kg/cm<sup>2</sup>.

In berries without any processing, the indicator was 0.21 and 0.24 kg/cm<sup>2</sup>. At the raspberry storage period end, the density of processed berries was in the range of 0.14-0.20 kg/cm<sup>2</sup> and 0.17-0.24 kg/cm<sup>2</sup>, depending on the storage method, which is 0.01-0.07 and 0.02-0.09 kg/cm<sup>2</sup> more than in control. The analysis of the research results showed that pre-processing of raspberries with chitosan solutions significantly affects tissue density preservation. As the concentration of the solution increased, the indicator decreased more slowly.

Ethyl alcohol is a strong solvent, and due to that, all biochemical processes are accelerated. Its formation in raspberries during storage occurs due to insufficient oxygen when the berries start anaerobic respiration [48].


The highest content of ethyl alcohol was observed in samples stored in a modified gas environment, 0.68-0.85% in processed berries, and 0.88% in the control (Figure 8).

In the samples that were stored with free access to air, the content of ethyl alcohol did not exceed 0.05-0.18%.



**Figure 8** The ethyl alcohol content in raspberries after storage (2018-2020): A – with free access to air; B – in a modified gas environment.


It was established that after fourteen days of storage, the accumulation of acetaldehyde was in raspberries (Figure 9). This indicates the creation of anaerobic conditions in the storage environment.



**Figure 9** The acetaldehyde content in raspberries after storage (2018-2020): A – with free access to air; B – in a modified gas environment.

It was found that the mass fraction of acetaldehyde was higher in samples stored in a modified gas environment. The indicator ranged from 0.11 to 0.15 mg/100 g when berries were stored with free access to air, and from 0.16 to 0.28 in MGE.

It was established that pre-processing of raspberries with chitosan solutions improved organoleptic properties (Figure 10), better-preserving colour, tissue density and taste.



**Figure 10** The organoleptic assessment of raspberries after storage (2018-2020): A – with free access to air; B – in a modified gas environment.

An organoleptic assessment is one of the most important indicators of product quality.

The potential buyer, first of all, pays attention to the appearance of the product, its color, aroma and consistency. The maintenance of the natural attraction of berries is a complex process, because during storage, a change in color, loss of elasticity and aroma is inevitable.

The accelerated rates of quality loss of raspberries were detected in the control sample. The sample with a 0.5% chitosan processing concentration was recognized as the most relevant for maintaining the organoleptic properties of raspberries.

Shine is a characteristic feature of the freshness of berry products. Its loss leads to the deterioration of the commercial qualities of the product, resulting in a decrease in the selling price.

Scientific papers [52] and [53] describe a wide range of organoleptic examinations of different types of berries immediately after harvesting, and also indicate the storage periods without the use of special solutions, but any studies of physical and chemical features were not conducted; it is unclear how the qualitative composition of raw materials changed during storage.

During the storage of raspberries, quick rates of shine loss were detected, particularly in the control sample (Table 1).

**Table 1** The changes in the raspberry shine degree before storage and during refrigerated storage (2018-2020).

Processing Concentration, %	Shine Degree, points					
	With free access to air			In a modified GE		
	Before storage	In the middle of the storage period (the 7th day)	At the storage period end (the 14th day)	Before storage	In the middle of the storage period (the 7th day)	At the storage period end (the 14th day)
Control	5	2	1	5	3	1
0.05	5	4	2	5	4	2
0.1	5	5	2	5	5	2
0.2	5	5	3	5	5	4
0.3	5	5	4	5	5	4
0.4	5	5	5	5	5	5
0.5	5	5	5	5	5	5
HIP <sub>05</sub>				0.2		


It was established that thanks to the film-forming properties of chitosan, the berries had a shiny glossy surface, significantly improving the raspberry's appearance. The obtained research results make it possible to recommend pre-processing raspberry with 0.4% and 0.5% concentration chitosan solutions.

Rapid losses of quality accompany the storage of berries in the conditions of trade transactions. This occurs due to increased physiological activity, particularly the high respiration intensity, which leads to the loss of nutritional substances [49].

During the storage of berries at a temperature of 20-22 °C, the quick deterioration of raspberry quality was observed. The criterion for the end of storage was the appearance of phytopathogenic damage on the surface of the berries. The berries were assessed according to the physical parameters. The investigations were conducted every day. It was found that the weight loss depended on both the storage period and the berry processing concentration. Scientific papers [50] and [51] describe the process of storing raspberries at temperatures above 22 °C, and the authors do not state the deterioration of quality indicators. Thus, the question arises about what preparation was used for the berries pre-processing; this information is not specified.

Thus, on the second day of non-refrigerated storage of berries, the mass loss ranged from 6.2 to 8.2% and from 6.4 to 8.4%, depending on the processing concentration, and from 8.3% to 8.6% in control. During the following days of the storage, the indicator increased and at the end, was 15.4-19.2% in the berries processed by the spraying method and 21.0% in control, and 15.1-19.0% in the samples processed by the immersion method and 20.9% in control.


The density of the berry tissues also decreased with the loss of mass. Thus, on the second day of the storage, the indicator decreased by 0.06-0.11 kg/cm<sup>2</sup> for both processing methods (Figure 11 and Figure 12).



**Figure 11** The change in the density of raspberries during non-refrigerated storage using the spraying processing method (2018-2020).

On the third day of storage, the density of berry tissues ranged from 0.15 to 0.20 kg/cm<sup>2</sup>. By the end of storage, the indicator decreased rapidly, and on the fifth day, it was 0.12 kg/cm<sup>2</sup> in samples with processing concentrations of 0.4 and 0.5% and 0.10 kg/cm<sup>2</sup> in the remaining samples.

The pH level of raspberries changed very quickly during storage in the conditions of trade transactions. The active acidity has decreased, which led to the deterioration of the quality of berries and the development of fungal diseases.



**Figure 12** The change in the density of raspberries during non-refrigerated storage with the immersion processing method (2018-2020).

On the second day of the storage, the indicator ranged from 3.2 to 3.5 for both processing methods. On the third day of storage, the pH level decreased to 3.6-4.4 for the spraying method and 3.8-4.2 for the immersion method. At the end of the storage period of the berries, the pH level ranged from 4.3 to 4.9 and from 4.4 to 4.8 in the processed samples and 5.0 in the control.

### CONCLUSION

It was established that the chitosan coating positively affects the maintenance of raspberries' quality indicators. It was detected that the processed berries had a loss of ascorbic acid of 1.5-3.9 times less than the control sample. A similar situation was observed with the mass fraction of sugars, which prevailed 0.9-2.5 times in the processed samples. According to the organoleptic indicators, raspberries without processing had an unattractive appearance, a softened consistency and a less pronounced aroma.

It was established that the mass loss of strawberries has increased daily regardless of the storage method. The strawberries' weight loss has a high inverse correlation dependence on the processing concentration. The mass loss has decreased with the increase of the chitosan percentage in the solution. The strawberries stored at a temperature of 20-22 °C for more than five days, both processed and unprocessed, are not advisable due to a significant loss of mass (15.4-21.0%). The ascorbic acid content accumulated during the vegetation quickly decreased on the second day of storage in all studied samples. The strawberries processed with chitosan solutions had lower vitamin C losses than the control. The



high inverse correlation dependence between the change in the ascorbic acid content and the chitosan solution concentration was established.

It was found that the change in the pH level of strawberries depends on the chitosan processing concentration, regardless of the storage and processing methods.

The tissue density decreased during the entire storage period and reached the value of 10 kg/cm<sup>2</sup>. It was investigated that strawberries processed with a chitosan solution of different concentrations had a higher density, contributing to long-term preservation.

The content of ethyl alcohol at the end of storage depended on the storage method. The highest values were detected when strawberries were stored in a modified gas environment. It was found that the strawberries pre-processed with a chitosan solution were suitable for preservation after seven days of storage. According to the physical, chemical, and organoleptic indicators, the preserves made from pre-processed berries fully complied with the requirements.

### REFERENCES

1. Cervantes, L., Martínez-Ferri, E., Soria, C., & Ariza, M. T. (2020). Bioavailability of phenolic compounds in strawberry, raspberry and blueberry: Insights for breeding programs. In *Food Bioscience* (Vol. 37, p. 100680). Elsevier BV. <https://doi.org/10.1016/j.fbio.2020.100680>
2. Asche, F., Straume, H., & Vårdal, E. (2021). Perish or prosper: Trade patterns for highly perishable seafood products. In *Agribusiness* (Vol. 37, Issue 4, pp. 876–890). Wiley. <https://doi.org/10.1002/agr.21704>
3. Rathod, N. B., Bangar, S. P., Šimat, V., & Ozogul, F. (2022). Chitosan and gelatine biopolymer-based active/biodegradable packaging for the preservation of fish and fishery products. In *International Journal of Food Science & Technology* (Vol. 58, Issue 2, pp. 854–861). Wiley. <https://doi.org/10.1111/ijfs.16038>
4. Wu, P., Xin, F., Xu, H., Chu, Y., Du, Y., Tian, H., & Zhu, B. (2021). Chitosan inhibits postharvest berry abscission of ‘Kyoho’ table grapes by affecting the structure of abscission zone, cell wall degrading enzymes and SO<sub>2</sub> permeation. In *Postharvest Biology and Technology* (Vol. 176, p. 111507). Elsevier BV. <https://doi.org/10.1016/j.postharvbio.2021.111507>
5. Eshghi, S., Karimi, R., Shiri, A., Karami, M., & Moradi, M. (2021). The novel edible coating based on chitosan and gum ghatti to improve the quality and safety of ‘Rishbaba’ table grape during cold storage. In *Journal of Food Measurement and Characterization* (Vol. 15, Issue 4, pp. 3683–3693). Springer Science and Business Media LLC. <https://doi.org/10.1007/s11694-021-00944-4>
6. Palamarchuk, I., Zozulyak, O., Mushtruk, M., Petrychenko, I., Slobodyanyuk, N., Domin, O., Udodov, S., Semenova, O., Karpovych, I., & Blishch, R. (2022). The intensification of dehydration process of pectin-containing raw materials. In *Potravinárstvo Slovak Journal of Food Sciences* (Vol. 16, pp. 15–26). HACCP Consulting. <https://doi.org/10.5219/1711>
7. Nguyen, V. T. B., Nguyen, D. H. H., & Nguyen, H. V. H. (2020). Combination effects of calcium chloride and nano-chitosan on the postharvest quality of strawberry (*Fragaria x ananassa* Duch.). In *Postharvest Biology and Technology* (Vol. 162, p. 111103). Elsevier BV. <https://doi.org/10.1016/j.postharvbio.2019.111103>
8. Rajestary, R., Landi, L., & Romanazzi, G. (2020). Chitosan and postharvest decay of fresh fruit: Meta-analysis of disease control and antimicrobial and eliciting activities. In *Comprehensive Reviews in Food Science and Food Safety* (Vol. 20, Issue 1, pp. 563–582). Wiley. <https://doi.org/10.1111/1541-4337.12672>
9. Celebi Sözüner, Z., Cevhertas, L., Nadeau, K., Akdis, M., & Akdis, C. A. (2020). Environmental factors in epithelial barrier dysfunction. In *Journal of Allergy and Clinical Immunology* (Vol. 145, Issue 6, pp. 1517–1528). Elsevier BV. <https://doi.org/10.1016/j.jaci.2020.04.024>
10. Ali, G., Sharma, M., Salama, E.-S., Ling, Z., & Li, X. (2022). Applications of chitin and chitosan as natural biopolymer: potential sources, pretreatments, and degradation pathways. In *Biomass Conversion and Biorefinery*. Springer Science and Business Media LLC. <https://doi.org/10.1007/s13399-022-02684-x>
11. Nikolaienko, M., & Bal-Prylypko, L. (2020). Development of an integrated food quality management system. In *Potravinárstvo Slovak Journal of Food Sciences* (Vol. 14, pp. 862–873). HACCP Consulting. <https://doi.org/10.5219/1434>
12. Madamsetty, V. S., Tavakol, S., Moghassemi, S., Dadashzadeh, A., Schneible, J. D., Fatemi, I., Shirvani, A., Zarrabi, A., Azedi, F., Dehshahri, A., Aghaei Afshar, A., Aghaabbasi, K., Pardakhty, A., Mohammadinejad, R., & Kesharwani, P. (2022). Chitosan: A versatile bio-platform for breast cancer theranostics. In *Journal of Controlled Release* (Vol. 341, pp. 733–752). Elsevier BV. <https://doi.org/10.1016/j.jconrel.2021.12.012>
13. DSTU 7179:2010 Fresh raspberries. Specifications. With correction. Quality management systems – Requirements.

14. DSTU 7205:2009. Fresh raspberries. Specifications. With correction. Quality management systems – Requirements.
15. DSTU 8102:2015 Canned foods. Fruit compotes for baby food. Specifications. Quality management systems – Requirements.
16. DSTU 4900:2007 Gems. General technical conditions. Quality management systems – Requirements.
17. John, O. D., Mouatt, P., Prasad, I., Xiao, Y., Panchal, S. K., & Brown, L. (2019). The edible native Australian fruit, Davidson's plum (*Davidsonia pruriens*), reduces symptoms in rats with diet-induced metabolic syndrome. In *Journal of Functional Foods* (Vol. 56, pp. 204–215). Elsevier BV. <https://doi.org/10.1016/j.jff.2019.03.018>
18. Lin, Y.-L., & Zheng, N.-Y. (2021). Torrefaction of fruit waste seed and shells for biofuel production with reduced CO<sub>2</sub> emission. In *Energy* (Vol. 225, p. 120226). Elsevier BV. <https://doi.org/10.1016/j.energy.2021.120226>
19. DSTU 8402:2015 Products of fruit and vegetable processing. Refractometric method of determining the content of soluble solids. Quality management systems – Requirements.
20. DSTU 4954:2008. Products of fruit and vegetable processing. Methods determination of sugars. Quality management systems – Requirements.
21. DSTU 4957:2008. Products of fruit and vegetable processing. Methods of determination of titrated acidity. Quality management systems – Requirements.
22. DSTU 6045:2008. Fruits, vegetables and processed products, canned meat and vegetables. Method for determining pH. Quality management systems – Requirements.
23. Shi, K., Liu, Z., Wang, J., Zhu, S., & Huang, D. (2019). Nitric oxide modulates sugar metabolism and maintains the quality of red raspberry during storage. In *Scientia Horticulturae* (Vol. 256, p. 108611). Elsevier BV. <https://doi.org/10.1016/j.scienta.2019.108611>
24. Yang, J., Cui, J., Chen, J., Yao, J., Hao, Y., Fan, Y., & Liu, Y. (2020). Evaluation of physicochemical properties in three raspberries (*Rubus idaeus*) at five ripening stages in northern China. In *Scientia Horticulturae* (Vol. 263, p. 109146). Elsevier BV. <https://doi.org/10.1016/j.scienta.2019.109146>
25. Gales, O., Rodemann, T., Jones, J., & Swarts, N. (2020). Application of near spectroscopy as an instantaneous and simultaneous prediction tool for anthocyanins and sugar in whole fresh raspberry. In *Journal of the Science of Food and Agriculture* (Vol. 101, Issue 6, pp. 2449–2454). Wiley. <https://doi.org/10.1002/jsfa.10869>
26. DSTU 4948:2008. Fruits, vegetables and their processing products. Methods of determination of nitrate content. Quality management systems – Requirements.
27. Vasylyshyna, O. V. (2020). Economic efficiency of preservation of cherry fruits for treatments by polysaccharidic compositions. In *Plant and Soil Science* (Vol. 11, Issue 2, pp. 74–80). National University of Life and Environmental Sciences of Ukraine. <https://doi.org/10.31548/agr2020.02.074>
28. Ivanova, I., Serdiuk, M., Tymoshchuk, T., Havryliuk, O., & Tonkha, V. (2022). Dynamics of the average fruit weight and the ratio of stone to pulp in the cherry fruit grown in the south of the steppe zone of Ukraine. In *PLANT AND SOIL SCIENCE* (Vol. 13, Issue 3). National University of Life and Environmental Sciences of Ukraine. [https://doi.org/10.31548/agr.13\(3\).2022.27-37](https://doi.org/10.31548/agr.13(3).2022.27-37)
29. Tereshchenko, N., Kovshun, L., & Bobunov, O. (2022). A hybrid technique for measuring the content of xenobiotics in wild and cultivated blueberries. In *PLANT AND SOIL SCIENCE* (Vol. 13, Issue 1, pp. 51–59). National University of Life and Environmental Sciences of Ukraine. [https://doi.org/10.31548/agr.13\(1\).2022.51-59](https://doi.org/10.31548/agr.13(1).2022.51-59)
30. Ni, J., Gao, J., Li, J., Yang, H., Hao, Z., & Han, Z. (2021). E-AlexNet: quality evaluation of strawberry based on machine learning. In *Journal of Food Measurement and Characterization* (Vol. 15, Issue 5, pp. 4530–4541). Springer Science and Business Media LLC. <https://doi.org/10.1007/s11694-021-01010-9>
31. Włodarska, K., Szulc, J., Khmelinskii, I., & Sikorska, E. (2019). Non-destructive determination of strawberry fruit and juice quality parameters using ultraviolet, visible, and near-infrared spectroscopy. In *Journal of the Science of Food and Agriculture* (Vol. 99, Issue 13, pp. 5953–5961). Wiley. <https://doi.org/10.1002/jsfa.9870>
32. Andrusyk, H. S., & Andrusyk, Y. Y. (2021). Qualitative characteristics of *Ribes Rubrum* L. varieties breeding at the department of horticulture named after prof. V.L. Symyrenko NULES of Ukraine. In *PLANT AND SOIL SCIENCE* (Vol. 12, Issue 1, pp. 120–128). National University of Life and Environmental Sciences of Ukraine. <https://doi.org/10.31548/agr2021.01.120>
33. Krstić, Đ., Milinčić, D. D., Kostić, A. Ž., Fotirić Akšić, M., Stanojević, S. P., Milojković-Opsenica, D., Pešić, M. B., & Trifković, J. (2022). Comprehensive electrophoretic profiling of proteins as a powerful tool for authenticity assessment of seeds of cultivated berry fruits. In *Food Chemistry* (Vol. 383, p. 132583). Elsevier BV. <https://doi.org/10.1016/j.foodchem.2022.132583>

34. Ivanova, I., Serdiuk, M., Malkina, V., Tonkha, O., Tsyz, O., Shkinder-Barmina, A., Verkholtantseva, V., Palianychka, N., Mushtruk, M., & Rozbytska, T. (2022). Factorial analysis of taste quality and technological properties of cherry fruits depending on weather factors. In *Potravinarstvo Slovak Journal of Food Sciences* (Vol. 16, pp. 341–355). HACCP Consulting. <https://doi.org/10.5219/1766>
35. Ivanova, I., Serdiuk, M., Malkina, V., Bandura, I., Kovalenko, I., Tymoshchuk, T., Tonkha, O., Tsyz, O., Mushtruk, M., & Omelian, A. (2021). The study of soluble solids content accumulation dynamics under the influence of weather factors in the fruits of cherries. In *Potravinarstvo Slovak Journal of Food Sciences* (Vol. 15, pp. 350–359). HACCP Consulting. <https://doi.org/10.5219/1554>
36. Mishra, P., Roger, J. M., Rutledge, D. N., & Woltering, E. (2020). SPORT pre-processing can improve near-infrared quality prediction models for fresh fruits and agro-materials. In *Postharvest Biology and Technology* (Vol. 168, p. 111271). Elsevier BV. <https://doi.org/10.1016/j.postharvbio.2020.111271>
37. Zheplinska, M., Mushtruk, M., Vasyliv, V., Slobodyanyuk, N., & Boyko, Y. (2021). The Main Parameters of the Physalis Convection Drying Process. In *Lecture Notes in Mechanical Engineering* (pp. 306–315). Springer International Publishing. [https://doi.org/10.1007/978-3-030-77823-1\\_31](https://doi.org/10.1007/978-3-030-77823-1_31)
38. Özdemir, İ. S., Bureau, S., Öztürk, B., Seyhan, F., & Aksoy, H. (2018). Effect of cultivar and season on the robustness of PLS models for soluble solid content prediction in apricots using FT-NIRS. In *Journal of Food Science and Technology* (Vol. 56, Issue 1, pp. 330–339). Springer Science and Business Media LLC. <https://doi.org/10.1007/s13197-018-3493-3>
39. Zheplinska, M., Mushtruk, M., Kos, T., Vasyliv, V., Kryzhova, Y., Mukoid, R., Bilko, M., Kuts, A., Kambulova, Y., & Gunko, S. (2020). The influence of cavitation effects on the purification processes of beet sugar production juices. In *Potravinarstvo Slovak Journal of Food Sciences* (Vol. 14, pp. 451–457). HACCP Consulting. <https://doi.org/10.5219/1284>
40. Dehghani, A., Bahlakeh, G., & Ramezanzadeh, B. (2019). A detailed electrochemical/theoretical exploration of the aqueous Chinese gooseberry fruit shell extract as a green and cheap corrosion inhibitor for mild steel in acidic solution. In *Journal of Molecular Liquids* (Vol. 282, pp. 366–384). Elsevier BV. <https://doi.org/10.1016/j.molliq.2019.03.011>
41. Kryzhova, Y., & Deyak, O. (2021). Investigation of the efficiency of beet syrup usage in products based on nutraceutical principles. In *Animal Science and Food Technology* (Vol. 12, Issue 2, pp. 40–47). National University of Life and Environmental Sciences of Ukraine. <https://doi.org/10.31548/animal2021.02.005>
42. Piechowiak, T., Skóra, B., & Balawejder, M. (2020). Ozone Treatment Induces Changes in Antioxidative Defense System in Blueberry Fruit During Storage. In *Food and Bioprocess Technology* (Vol. 13, Issue 7, pp. 1240–1245). Springer Science and Business Media LLC. <https://doi.org/10.1007/s11947-020-02450-9>
43. Burova, Z. A., Ivanov, S. O., Roman, T. O., Vasyliv, V. P., Zheplinska, M. M., Mushtruk, M. M., Palamarchuk, I. P., Sarana, V. V., & Gudzenko, M. M. (2021). Research on thermophysical characteristics of food products. In *Animal Science and Food Technology* (Vol. 12, Issue 3, pp. 18–35). National University of Life and Environmental Sciences of Ukraine. <https://doi.org/10.31548/animal2021.03.002>
44. Jiang, Y., Luo, T., Tang, Y., Chen, S., Ni, H., Chen, Q., Song, X., Bao, Y., Deng, Z., & Wang, J. (2022). Isolation of a novel characterized *Issatchenkia terricola* from red raspberry fruits on the degradation of citric acid and enrichment of flavonoid and volatile profiles in fermented red raspberry juice. In *Food Science and Human Wellness* (Vol. 11, Issue 4, pp. 1018–1027). Elsevier BV. <https://doi.org/10.1016/j.fshw.2022.03.029>
45. Gu, X., Xue, L., Lu, L., Xiao, J., Song, G., Xie, M., & Zhang, H. (2020). Melatonin Enhances the Waterlogging Tolerance of *Prunus persica* by Modulating Antioxidant Metabolism and Anaerobic Respiration. In *Journal of Plant Growth Regulation* (Vol. 40, Issue 5, pp. 2178–2190). Springer Science and Business Media LLC. <https://doi.org/10.1007/s00344-020-10263-5>
46. Zheplinska, M., Mushtruk, M., Vasyliv, V., Kuts, A., Slobodyanyuk, N., Bal-Prylypko, L., Nikolaenko, M., Kokhan, O., Reznichenko, Y., & Salavor, O. (2021). The micronutrient profile of medicinal plant extracts. In *Potravinarstvo Slovak Journal of Food Sciences* (Vol. 15, pp. 528–535). HACCP Consulting. <https://doi.org/10.5219/1553>
47. Palamarchuk, I., Palamarchuk, V., Sarana, V., Heipel, Y., & Borodych, B. (2022). Technical and economic substantiation of the process of semi-fluidisation treatment of fruit and berry products. In *Animal Science and Food Technology* (Vol. 13, Issue 2, pp. 35–43). National University of Life and Environmental Sciences of Ukraine. [https://doi.org/10.31548/animal.13\(2\).2022.35-43](https://doi.org/10.31548/animal.13(2).2022.35-43)
48. Joseph, M., Van Beers, R., Postelmans, A., Nicolai, B., & Saeys, W. (2021). Exploring oxygen diffusion and respiration in pome fruit using non-destructive gas in scattering media absorption spectroscopy. In *Postharvest Biology and Technology* (Vol. 173, p. 111405). Elsevier BV. <https://doi.org/10.1016/j.postharvbio.2020.111405>

49. Piechowiak, T., Antos, P., Kosowski, P., Skrobacz, K., Józefczyk, R., & Balawejder, M. (2019). Impact of ozonation process on the microbiological and antioxidant status of raspberry (*Rubus idaeus* L.) fruit during storage at room temperature. In *Agricultural and Food Science* (Vol. 28, Issue 1). Agricultural and Food Science. <https://doi.org/10.23986/afsci.70291>
50. Piechowiak, T., & Balawejder, M. (2019). Impact of ozonation process on the level of selected oxidative stress markers in raspberries stored at room temperature. In *Food Chemistry* (Vol. 298, p. 125093). Elsevier BV. <https://doi.org/10.1016/j.foodchem.2019.125093>
51. Kos, T., Kuznietsova, I., Sheiko, T., Khomichak, L., Kambulova, Y., Bal-Prylypko, L., Vasylyv, V., Nikolaenko, M., Bondar, M., & Babych, I. (2021). An improved method for determining the mass fraction of calcium carbonate in the carbonate bedrock. In *Potravinarstvo Slovak Journal of Food Sciences* (Vol. 15, pp. 877–890). HACCP Consulting. <https://doi.org/10.5219/1591>
52. Palamarchuk, I., Kiurchev, S., Verkhohantseva, V., Borodych, B., & Lebska, T. (2022). Justification of power parameters of the process of semi-fluidisation freezing of fruit and berry products. In *Animal Science and Food Technology* (Vol. 13, Issue 1, pp. 39–46). National University of Life and Environmental Sciences of Ukraine. [https://doi.org/10.31548/animal.13\(1\).2022.39-46](https://doi.org/10.31548/animal.13(1).2022.39-46)
53. Covaliov, E., Grosu, C., Popovici, V., Capcanari, T., Siminiuc, R., & Resitca, V. (2021). Impact of sea buckthorn berries (*Hippophae rhamnoides*) on yoghurt biological value and quality. In *The Annals of the University Dunarea de Jos of Galati Fascicle VI – Food Technology* (Vol. 45, Issue 2, pp. 62–76). Universitatea Dunarea de Jos din Galati. <https://doi.org/10.35219/foodtechnology.2021.2.05>

### Funds:

This research received no external funding.

### Acknowledgments:

We would like to thank you to Anastasiia Blahopoluchna.

### Conflict of Interest:

No potential conflict of interest was reported by the author(s).


### Ethical Statement:

This article does not contain any studies that would require an ethical statement.

### Contact Address:

**Anastasiia Blahopoluchna**, Pavlo Tychyna Uman State Pedagogical University; Institute of Economics and Business Education, Department of Technologies and Organization of Tourism and Hotel and Restaurant Business. Sadova str., 2, Uman, Cherkasy region, 20300, Ukraine,

E-mail: [a.blagopoluchna1995@gmail.com](mailto:a.blagopoluchna1995@gmail.com)

 ORCID: <https://orcid.org/orcid=0000-0001-5897-0120>

\***Mikhailo Mushtruk**, National University of Life and Environmental Sciences of Ukraine, Faculty of Food Technology and Quality Control of Agricultural Products, Department of Processes and Equipment for Processing of Agricultural Production, Heroes of Defense Str., 12 B, Kyiv, 03040, Ukraine,

E-mail: [mixej.1984@ukr.net](mailto:mixej.1984@ukr.net)

 ORCID: <https://orcid.org/0000-0002-3646-1226>

**Nataliia Slobodyanyuk**, National University of Life and Environmental Sciences of Ukraine, Faculty of Food Technology and Quality Control of Agricultural Products Department of technology of meat, fish and marine products, Polkovnyka Potiekhina Str., 16, Kyiv, 03040, Ukraine,

E-mail: [slob2210@ukr.net](mailto:slob2210@ukr.net)

 ORCID: <https://orcid.org/0000-0002-7724-2919>

**Nelia Liakhovska**, Uman National University of Horticulture., Department of Biology, Instytutska Str., 1 (building 4), Uman, 20305, Cherkasy Region, Ukraine,

E-mail: [lyakhovska@i.ua](mailto:lyakhovska@i.ua)

 ORCID: <https://orcid.org/0009-0001-0995-8966>

**Vladyslav Parakhnenko**, Pavlo Tychyna Uman State Pedagogical University; Institute of Economics and Business Education, Department of chemistry, ecology and methods of their teaching, Sadova str., 2, Uman, 20300, Cherkasy region, Ukraine,

E-mail: [vladparachenko@ukr.net](mailto:vladparachenko@ukr.net)

 ORCID: <https://orcid.org/0000-0002-4312-6194>

**Sergiy Udodov**, National University of Food Technologies, Department of Machines and Apparatuses for Food and Pharmaceutical Productions, Volodymyrska Str. 68, 01601 Kyiv, Ukraine,

E-mail: [kiypivo@gmail.com](mailto:kiypivo@gmail.com)

 ORCID: <https://orcid.org/0000-0001-6149-9974>


**Inna Karpovych**, National University of Food Technologies, Department of Sugar Technology and Water Preparation, Volodymyrska Str. 68, 01601 Kyiv, Ukraine,

E-mail: [inkarp@ukr.net](mailto:inkarp@ukr.net)

 ORCID: <https://orcid.org/0000-0002-5912-7466>

**Olena Ochkolyas**, The National University of Life and Environmental Sciences of Ukraine, Faculty of Food Technology and Quality Control of Agricultural Products, Department of technology of meat, fish and marine products, Polkovnika Potekhina, Str., 16, Kyiv, 02163, Ukraine,

E-mail: [ochkolyas@nubip.edu.ua](mailto:ochkolyas@nubip.edu.ua)

 ORCID: <https://orcid.org/0000-0002-8483-578X>

**Alina Omelian**, National University of Life and Environmental Sciences of Ukraine, Faculty of Food Technology and Quality Control of Agricultural Products, Department of Technologies of Meat, Fish and Seafoods, Heroiv Oborony Str., 15, 03040, Kyiv, Ukraine,

E-mail: [alina\\_omelian@nubip.edu.ua](mailto:alina_omelian@nubip.edu.ua)

 ORCID: <https://orcid.org/0000-0001-9004-5250>

**Gennadiy Rzhevsky**, National University of Life and Environmental Sciences of Ukraine, Faculty of Humanities and Pedagogy Department of Psychology, lane Agricultural 4b, Kyiv, 03041, Ukraine,

E-mail: [manpower44444@gmail.com](mailto:manpower44444@gmail.com)

 ORCID: <https://orcid.org/0000-0003-0515-1687>

Corresponding author: \*

© 2023 Authors. Published by HACCP Consulting in [www.potravinarstvo.com](http://www.potravinarstvo.com) the official website of the *Potravinarstvo Slovak Journal of Food Sciences*, owned and operated by the HACCP Consulting s.r.o., Slovakia, European Union [www.haccp.sk](http://www.haccp.sk). The publisher cooperate with the SLP London, UK, [www.slplondon.org](http://www.slplondon.org) the scientific literature publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License CC BY-NC-ND 4.0 <https://creativecommons.org/licenses/by-nc-nd/4.0/>, which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.